

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 9/16, 9/54, 9/62	A1	(11) International Publication Number: WO 00/03696 (43) International Publication Date: 27 January 2000 (27.01.00)
(21) International Application Number: PCT/US98/16323 (22) International Filing Date: 6 August 1998 (06.08.98) (30) Priority Data: 09/118,418 17 July 1998 (17.07.98) US (71) Applicant: BRISTOL-MYERS SQUIBB COMPANY [US/US]; P.O. Box 4000, Princeton, NJ 08543-4000 (US). (72) Inventors: ULLAH, Ismat; 2 Mockingbird Court, Cranbury, NJ 08512 (US). WILEY, Gary, J.; 72 Tuscany Drive, Jackson, NJ 08527 (US). (74) Agents: MARENBERG, Barry, J. et al.; Bristol-Myers Squibb Company, P.O. Box 4000, Princeton, NJ 08543-4000 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: ENTERIC COATED PHARMACEUTICAL TABLET AND METHOD OF MANUFACTURING (57) Abstract A high drug load enteric coated pharmaceutical composition is provided which includes a core in the form of a tablet and which is comprised of a medicament which is sensitive to a low pH environment of less than 3, such as ddI, and having an enteric coating formed of methacrylic acid copolymer and a plasticizer. The tablets may be of varying sized and may be orally ingested individually or a plurality of tablets sufficient to attain a desired dosage may be encapsulated in a dissolvable capsule. The tablets have excellent resistance to disintegration at pH less than 3 but have excellent drug release properties at pH greater than 4.5. A novel method of making said pharmaceutical composition is also disclosed.		

LEDIGLICH ZUR INFORMATION

Codes zur Identifizierung von PCT-Vertragsstaaten auf den Kopfbögen der Schriften, die internationale Anmeldungen gemäss dem PCT veröffentlichen.

AL	Albanien	ES	Spanien	LS	Lesotho	SI	Slowenien
AM	Armenien	FI	Finnland	LT	Litauen	SK	Slowakei
AT	Österreich	FR	Frankreich	LU	Luxemburg	SN	Senegal
AU	Australien	GA	Gabun	LV	Lettland	SZ	Swasiland
AZ	Aserbaidschan	GB	Vereinigtes Königreich	MC	Monaco	TD	Tschad
BA	Bosnien-Herzegowina	GE	Georgien	MD	Republik Moldau	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagaskar	TJ	Tadschikistan
BE	Belgien	GN	Guinea	MK	Die ehemalige jugoslawische Republik Mazedonien	TM	Turkmenistan
BF	Burkina Faso	GR	Griechenland	ML	Mali	TR	Türkei
BG	Bulgarien	HU	Ungarn	MN	Mongolei	TT	Trinidad und Tobago
BJ	Benin	IE	Irland	MR	Mauretanien	UA	Ukraine
BR	Brasilien	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Island	MX	Mexiko	US	Vereinigte Staaten von Amerika
CA	Kanada	IT	Italien	NE	Niger	UZ	Usbekistan
CF	Zentralafrikanische Republik	JP	Japan	NL	Niederlande	VN	Vietnam
CG	Kongo	KE	Kenia	NO	Norwegen	YU	Jugoslawien
CH	Schweiz	KG	Kirgisistan	NZ	Neuseeland	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Demokratische Volksrepublik Korea	PL	Polen		
CM	Kamerun	KR	Republik Korea	PT	Portugal		
CN	China	KZ	Kasachstan	RO	Rumänien		
CU	Kuba	LC	St. Lucia	RU	Russische Föderation		
CZ	Tschechische Republik	LI	Liechtenstein	SD	Sudan		
DE	Deutschland	LK	Sri Lanka	SE	Schweden		
DK	Dänemark	LR	Liberia	SG	Singapur		
EE	Estland						

ENTERIC COATED PHARMACEUTICAL TABLET AND METHOD OF MANUFACTURING

Brief Description of the Invention

5 The present invention is directed to an enteric-coated pharmaceutical composition in the form of a tablet which comprises an acid labile high drug load medicament, such as ddl, which is sensitive to a low pH environment of less than 3, and which includes an enteric coating such as Eudragit L-30-D 55 and a plasticizer, but which does not require a subcoat. The tablets have excellent resistance to disintegration at pH less than 3 but have excellent drug release properties at pH greater than 4.5. A novel method of making said pharmaceutical composition is also disclosed.

15 **Background of the Invention**

Enteric coatings have been used for many years to arrest the release of the drug from orally ingestible dosage forms. Depending upon the composition and/or thickness, the enteric coatings are resistant to stomach acid for required periods of time before they begin to disintegrate and permit slow release of the drug in the lower stomach or upper part of the small intestines. Examples of some enteric coatings are disclosed in U.S. Patent No. 5,225,202 which is incorporated by reference fully herein. As set forth in U.S. Patent No. 5,225,202, some examples of coating previously employed are beeswax and glyceryl monostearate; beeswax, shellac and cellulose; and cetyl alcohol, mastic and shellac, as well as shellac and stearic acid (U.S. Pat. No. 2,809,918); polyvinyl acetate and ethyl cellulose (U.S. Pat. No. 3,835,221); and neutral copolymer of polymethacrylic acid esters (Eudragit L30D) (F.W. Goodhart et al., Pharm. Tech., pp. 64-71, April 1984); copolymers of methacrylic acid and methacrylic acid methylester (Eudragits), or a neutral copolymer of polymethacrylic acid esters containing metallic stearates (Mehta et al., U.S. Pat. Nos. 4,728,512 and 4,794,001).

Most enteric coating polymers begin to become soluble at pH 5.5 and above, with maximum solubility rates at pHs greater than 6.5.

35 Numerous enteric coated and/or extended release pharmaceutical compositions and the methods of making these compositions have been disclosed in the art. Prior art compositions, however, often comprise

numerous extra ingredients in addition to the medicaments, such as fillers, buffering agents, binders and wetting agents, all of which add to the bulk of the composition and reduce the amount of active medicament which can be contained in the composition. The processes for preparing these

5 aforementioned pharmaceutical compositions require multiple time consuming steps, including subcoating and outer coating steps. Furthermore, many of these pharmaceutical compositions are intended for delivery in the lower GI tract, i.e. in the colon, as opposed to the upper intestines, i.e. the duodenum of the small intestine.

10 U.S. Pat. No. 5,225,202 discloses enteric coated pharmaceutical compositions utilizing neutralized hydroxypropyl methylcellulose phthalate polymer (HPMCP) coating. The pharmaceutical compositions disclosed comprise an acid labile medicament core, a disintegrant, one or more buffering agents to provide added gastric protection in addition to the enteric coating, as well as the enteric coating and a plasticizer. The pharmaceutical composition may also include one or more lactose, sugar or starch fillers. According to the invention disclosed in this reference, when the core includes a drug which is incompatible with the enteric coating layer, an additional subcoat layer which acts as a physical barrier between the core and outer enteric coating layer is employed to prevent interaction of the acid labile drug and the acidic enteric coat. The HPMCP enteric coating starts its dissolution process at pH 5.0. The process of preparing this pharmaceutical composition requires numerous coating steps to apply the subcoat and then the enteric coat.

25 U.S. Pat. No. 5,026,560 discloses a pharmaceutical composition and method of making said pharmaceutical composition, wherein the pharmaceutical composition comprises a Nonpareil seed core produced by coating sucrose with corn starch, spraying the core with an aqueous binder in a solution of water or ethanol and with a spraying powder containing a drug and low substituted hydroxypropylcellulose, followed by the application of an enteric coating.

30 U.S. Pat. No. 4,524,060 recites a slow release pharmaceutical composition which provides a sustained release composition for treating hypertensive patients, and which comprises a mixture of micronized indoramin or a pharmaceutically acceptable salt thereof, a water-channeling agent, a wetting agent, a disintegrant, the mixture being in the form of a non-


compressed pellet and having an enteric coat or sustained release coat permeable to gastrointestinal juices.

U.S. Pat. No. 5,536,507 is directed to a pharmaceutical composition having a delayed release coating or enteric coatings wherein
5 the active agent in the composition is intended for release of a predominant amount of the drug at a point near the inlet to or within the large intestine and at a pH of approximately 6.4-7.0.

Pharmaceutical compositions which include a medicament which is unstable in an acidic environment such as the stomach and which is not
10 adequately buffered, will require an enteric protective coating to prevent release of such medicament prior to reaching the intestines.

ddl, (also known as didanosine or 2',3'-dideoxyinosine, and marketed by Bristol-Myers Squibb Co. under the brand name Videx®), is an acid labile drug which has the formula

15



and which has been shown to be effective in the treatment of patients with the HIV virus which causes AIDS. The composition and method of inhibiting HIV replication with 2',3'-dideoxyinosine have been reported. See U.S.
20 Patent Nos. 4,861,759, 5,254,539 and 5,616,566, which are incorporated by reference herein. More recently, Videx® has become widely used as a component of the new therapeutic cocktails used to treat AIDS. It is also an acid labile medicament sensitive to a low pH environment and will degrade in the stomach.

25 Videx® is generally available in a variety of oral dosages, including Chewable/Dispersible Buffered Tablets in strengths of 25, 50, 100 or 150 mg of didanosine. Each tablet is buffered with calcium carbonate and magnesium hydroxide. Videx® tablets also contain aspartame, sorbitol, microcrystalline cellulose, Polyplasdone®, mandarin-orange flavor, and
30 magnesium stearate. Videx® Buffered Powder for Oral Solution is supplied

for oral administration in single-dose packets containing 100, 167 or 250 mg of didanosine. Packets of each product strength also contain a citrate-phosphate buffer (composed of dibasic sodium phosphate, sodium citrate, and citric acid) and sucrose. A Videx® Pediatric Powder for Oral Solution is
5 also available and which is supplied for oral administration in 4- or 8-ounce glass bottles containing 2 or 4 grams of didanosine respectively, and is to be mixed with commercial antacid before oral ingestion.

With particular emphasis on the tablets, whether ingested alone or as part of a combination ("cocktail") therapy regimen, the current
10 chewable/dispersible buffered tablets are not conducive from a patient ease of use standpoint. Whereas the other products which are a part of the AIDS therapeutic cocktail are capsules or tablets and easily swallowed, the Videx® (referred to herein as "ddl") Chewable/Dispersible Buffered Tablets must be thoroughly chewed, manually crushed, or uniformly dispersed in
15 water before administration. Because ddl degrades rapidly at acidic pH, ddl, in its chewable/dispersible form and its buffered powder for oral solution, contains buffering agents and is administered with antacids in the pediatric powder form. However, the presence of the large quantities of antacid components in the formulation can lead to significant GI imbalance
20 as noted by severe diarrhea. Many patients also complain about chewing the large ddl tablets (dose = 2 tablets of 2.1 g each), the taste of the ddl or the time required to disperse the tablets and the volume of fluid (4 oz) required for the dose. All these factors, coupled with the fact that other nucleoside analog drugs are marketed in a more convenient dosage
25 presentation (i.e. capsule or smaller tablets), necessitate the development of an innovative dosage form of ddl which is easy to swallow and does not cause discomforting side effects.

Accordingly, there is provided a tablet comprising a medicament core and having a coating which prevents release of the medicament in the
30 stomach and allows for release of the drug in the small intestine thereby eliminating the need for an antacid which may cause GI imbalance upon chronic use. Thus, pharmaceutical compositions which include a medicament which is unstable in an acid environment such as the stomach will require such a protective coating to prevent release of such medicament
35 prior to reaching the intestines.

Description of the Invention

In accordance with the present invention, an enteric coated, high drug load pharmaceutical composition, and a method of making said pharmaceutical composition, is provided which includes a medicament which may degrade in a low pH environment but which is protected from doing so by the enteric coating. The pharmaceutical composition of the invention, which is advantageously in the form of tablets, includes a core which comprises a medicament, such as ddl, which is sensitive to a low pH environment and optionally a binder or filler, a disintegrant or swelling agent, and a lubricant. The core further comprises an enteric coating surrounding the core which includes a methacrylic acid copolymer and a plasticizer.

The novel enteric coated pharmaceutical of the invention will provide for protection of the medicament or therapeutically active agent, such as ddl, at pH's less than 3 (such as found in the stomach) but will permit drug release at a pH of 4.5 or higher (such as found in the upper intestines).

Accordingly, the pharmaceutical composition of the invention will usually include drugs which are chemically unstable in acidic environments. The pharmaceutical composition of the invention provides excellent protection in very acidic environments (pH <3) while not delaying the rapid release in regions of pH greater than 4, whether this be the upper intestine or the duodenum.

Most of the enteric coating materials known in the art are acidic in nature and hence may cause chemical instability when in contact with acid labile ingredients. This is especially true under high temperature and humid conditions experienced during an aqueous coating process. To minimize this acid caused instability, a protective coat or subcoat is usually applied between the particles, beadlets, pellets, tablets, etc., and the enteric coat. This protective coat physically separates the acid labile drug from the acidic enteric coat, and hence improves stability of the formulation. The process of applying such a subcoat, however, often entails multiple burdensome and time-consuming steps. Furthermore, the subcoat can cause a delay in drug release.

A process is thus described by which tablets, beadlets, pellets, and/or particles containing acid labile drugs can be successfully aqueous

enteric coated without application of the protective coat or subcoat. This process involves raising the pH of the enteric coating suspension solution by using alkalizing agents. The pH of the coating suspension is raised below the point where enteric integrity of the polymer could be lost. The
5 process may also involve the inclusion of binders, such as sodium carboxymethylcellulose, fillers, such as microcrystalline cellulose, disintegrants, such as sodium starch glycolate, and other excipients, such as magnesium oxide, which are relatively alkaline in nature, in the formulations of cores intended for enteric coating. Raising the pH of the
10 coating suspension provides a more stable composition for the acid labile drug in the core. As a result, there is no incompatibility and no need for a protective subcoat between the acid labile drug and the acidic enteric coat. This process not only eliminates the costly additional subcoating step, but allows quicker release of the drug since the added subcoat layer delays
15 drug release.

The process of the present invention illustrates the preparation of high (up to 99.5%) potency (uncoated) tablets, for acid labile drugs, such as ddl, using an aqueous process. No specialized equipment is required as conventional blending, compacting, tableting, and coating equipment was
20 found to be adequate for tablet formation and coating.

In the digestive tract, the coated tablets pass through the stomach first. The transit time for the stomach is approximately two hours and the pH of this region is approximately 1 to 3. The enteric coating component allows the medicament core to remain substantially intact and thus prevents the
25 pharmacologically active substance from being released in this region or the acid from penetrating through to the tablet core. The tablets then pass through the small intestine wherein the majority of the enteric coating component will dissolve and release the pharmacologically active substance therein. In normal flow direction therethrough, the small intestine
30 consists of the duodenum, jejunum and ileum. Transit time through the small intestine is approximately 2-4 hours and the pH of these regions is approximately 5 to approximately 7.2.

As used herein "enteric coating", is a polymer material or materials which encase the medicament core. The polymeric enteric coating material
35 in the present invention does not contain any active compound, i.e. any therapeutically active agent, of the present invention. Preferably, a substantial amount or all of the enteric polymer coating material is

dissolved before the medicament or therapeutically active agent is released from the dosage form, so as to achieve delayed dissolution of the medicament core. A suitable pH-sensitive polymer is one which will dissolve in intestinal juices at the higher pH levels (pH greater than 4.5),
5 such as within the small intestine and therefore permit release of the pharmacologically active substance in the regions of the small intestine and not in the upper portion of the GI tract, such as the stomach.

The polymer coating material is selected such that the therapeutically active agent will be released when the dosage form reaches
10 the small intestine or a region in which the pH is greater than pH 4.5. Preferred coating pH-sensitive materials, which remain intact in the lower pH environs of the stomach, but which disintegrate or dissolve at the pH commonly found in the small intestine of the patient. The enteric polymer coating material begins to dissolve in an aqueous solution at pH between
15 about 4.5 to about 5.5. The pH-solubility behavior of the enteric polymers of the present invention are such that significant dissolution of the enteric polymer coating will not occur until the dosage form has emptied from the stomach. The pH of the small intestine gradually increases from about 4.5 to about 6.5 in the duodenal bulb to about 7.2 in the distal portions of the
20 small intestine (ileum). In order to provide predictable dissolution corresponding to the small intestine transit time of about 3 hours and permit reproducible release therein, the coating should begin to dissolve within the pH range of the duodenum and continue to dissolve at the pH range within the small intestine. Therefore, the amount of enteric polymer coating should
25 be such that it is substantially dissolved during the approximate three hour transit time within the small intestine.

The pharmaceutical medicament present in the core will be an acid labile drug such as ddl, pravastatin, erythromycin, digoxin, pancreatin, ddA, ddC, and the like. The present invention is not limited to these drugs
30 and other drugs may be used as well. The invention is particularly adapted to pharmaceutical compositions, such as tablets, which contain ddl as the medicament. ddl will be present in an amount of about up to about 95% of the composition in the coated tablets.

One or more binders or fillers may be present in the core.
35 Microcrystalline cellulose (PH-101) is the preferred binder most suitable for use herein. Examples of other binders which may be used include sodium carboxymethylcellulose Avicel™ PH101, Avicel™ RC 591, Avicel™ CL-611,

(FMC Corp), Ceolus™ (FMC Corp.), ProSolv™ (Edward Mendell Co.) Methocel™ E-5 (Dow Corp.), Starch 1500 (Colorcon, Ltd.), Hydroxypropyl Methylcellulose (HPMC) (Shin-Etsu Chemical Co., Ltd.), Polyvinylpyrrolidone, Potassium Alginate and Sodium Alginate.

5 The core of the composition of the invention may also include one or more disintegrants or swelling agents, such as sodium starch glycolate marketed under the trademark EXPLOTAB (Edward Mendell Co.), Ac-Di-Sol (cross-linked sodium carboxymethylcellulose) (FMC Corp), croscarmellose sodium, corn starch, or cross linked polyvinylpyrrolidone. A lubricant such
10 as magnesium stearate, may also be used in the preparation of the uncoated tablet, specifically as a lubricant for the compaction and tableting process.

 The core employed in the pharmaceutical composition of the invention will be formed of a tablet, preferably a round, biconvex tablet,
15 approximately 3/16 of an inch. The invention is not, however, limited in the size of the tablet and tablets of varying sizes may be made. Smaller sized tablets are advantageous, however, since they pass through the stomach with more ease than larger sized tablets. Experimentation has shown that the tablet of the present invention having core comprising ddi as the
20 medicament, has the same bioavailability as the beadlet disclosed in co-pending U.S. Application No. 09/083,597, which was filed on May 22, 1998. Depending upon the size of the tablets, they may be ingested individually, or a plurality of tablets sufficient to attain a particular dosage may be encapsulated in a dissolvable capsule.

25 In an alternative embodiment of the present invention, the core can be prepared from a wet granulation process, using any of the wet granulation binders (if necessary) commonly used in the art, such as pregelatinized starch, polyvinylpyrrolidone, HPMC sodium carboxymethylcellulose, potassium or sodium alginate. The wet granulation
30 process comprises the steps of preparing granules suitable for tableting by blending a mixture comprising the medicament, a binder, and optionally, a disintegrant and filler; adding a predetermined amount of water or granulation solvent to form a wet mass blend; sizing the wet mass blend into granules to aid drying; drying the wet granules to remove excess
35 moisture; sizing the dried granules into granules suitable for tableting, and adding lubricant, one or more fillers, one or more dry binders, optionally a disintegrant, and other excipients necessary for tableting the granules.

The enteric coating according to the present invention will include methacrylic acid copolymer, a plasticizer, and a sufficient quantity of NaOH to adjust the pH of the suspension. Other alkalizing agents, such as potassium hydroxide, calcium carbonate, sodium carboxymethylcellulose, magnesium oxide, and magnesium hydroxide can also be used.

In forming the enteric coated pharmaceutical composition of the invention, an enteric coating solution of Eudragit L-30-D 55 will be employed. Eudragit L-30-D 55 is an aqueous acrylic resin dispersion, an anionic copolymer derived from methacrylic acid and ethyl acrylate with a ratio of free carboxyl groups to the ester of approximately 1:1, and a mean molecular weight of approximately 250,000, is supplied as an aqueous dispersion containing 30% w/w of dry lacquer substance, and is marketed by Rohm-Pharma Co., Germany. As an aqueous-based coating, no dangerous or environmentally harmful organic solvents are utilized.

Although Eudragit L-30-D-55 is the preferred coating polymer, the invention is not limited in this respect and other enteric coating polymers known in the art, such as hydroxypropyl methylcellulose phthalate HP50 (HPMCP-HP50) (USP/NF 220824), HP55 (HPMCP-HP55)(USP/NF type 200731) and HP55S available from Shin Etsu Chemical, Coateric™ (polyvinyl acetate phthalate)(Colorcon Ltd.), Sureteric™ (polyvinyl acetate phthalate)(Colorcon, Ltd.), or Aquateric™ (cellulose acetate phthalate)(FMC Corp.), and the like may be employed

The enteric coating will also preferably contain a plasticizer which is preferably diethyl phthalate, although the invention is not limited in this respect and other plasticizers may be used such as triethyl citrate (Citroflex- 2), triacetin, tributyl sebecate, or polyethylene glycol.

The enteric coating employed in the present invention is substantially easier to process than previously reported coating systems, and is especially advantageous for coating small diameter, low mass particles (tablets) with minimal processing problems (sticking/picking) without the need for organic solvents.

In general, where the core includes a drug which is incompatible with the enteric coating layer, a subcoat layer which may be comprised of one or more film-formers or plasticizers, and which acts as a physical barrier between the core and the outer enteric coating layer will be employed. However, unlike previously reported coatings such as that disclosed in U.S. Pat. No. 5,225,202, the novel pharmaceutical composition

of the present invention, as a result of the novel process utilized in making the composition of the present invention and the pH adjustment of the coating, does not require a subcoat since the need for such an insulating layer is eliminated by raising the pH of the aqueous coating suspension.

- 5 Since the coating is designed to breakdown at pH 5.5, the enteric coating applied at pH 5 permits relatively rapid breakdown in the intestine as only a small amount of additional alkalinity is required to bring the pH to 5.5.

A preferred formulation for preparing a 50 mg uncoated tablet is set out below.

10

Material		Amount (mg) per Tablet
<u>TABLET CORE</u>		
Drug (didanosine)		50.00
15 Microcrystalline Cellulose		17.00
Na Starch Glycolate		2.10
Magnesium Stearate (for compaction)		0.60
Magnesium Stearate (for tableting)		0.30
20 Uncoated Tablet Net Weight		<u>70.00</u>

A preferred formulation for the preparation of an enteric film coating suspension to coat the uncoated 50 mg tablets is set out below.

25

Material		Amount (g) per 100 g
<u>COATING</u>		
Eudragit L-30-D 55		66.67
Diethyl Phthalate		3.00
30 Purified Water		qs
(pH adjusted to 5 ± 0.1 with NaOH solution)		

- The percentage range of the ingredients in the above formulations
 35 for the uncoated tablet and the enteric film coating is set forth in the following chart:

40

	Material	% (range)
	<u>CORE</u>	
5	Drug (didanosine)	1 - 100
	Microcrystalline Cellulose	0 - 40
	Na Starch Glycolate	0 - 6
	Magnesium Stearate	0 - 3
10	<u>COATING</u>	
	Eudragit L-30-D 55	2 - 30
	Diethyl Phthalate	0.5 - 6.0

15 The enteric coated pharmaceutical composition in the form of
tablets may be prepared by a process which comprises the steps of mixing
an acid labile medicament, a binder/filler, such as microcrystalline
cellulose, a disintegrant, such as sodium starch glycolate, and a first portion
20 type blender, to prepare a dry blend. The blend is then screened and placed
back in the blender for a second blending. The resulting blend is slugged
or compacted and then sized to form small granules. A second portion of
magnesium stearate lubricant for tableting is then calculated and blended
in the tumbling type blender with the screened granules. The resulting
25 blend is then formed into tablets (uncoated) having a desired weight and
hardness.

 The tablets may then be coated with an enteric film coating
suspension comprising Eudragit L-30-D 55 and plasticizer (diethyl
phthalate), using a fluid bed coating apparatus with top spray mode, such
30 as an Aeromatic STREA-1 table top unit, and then dried. During preparation
of the film coating suspension, a NaOH solution is added to the suspension
until a pH of 5.0 ± 0.1 is obtained. Adjustment of the enteric film coating
suspension to pH 5 eliminates the need for a subcoat or insulating layer.
The advantage here is that an enteric coating at pH 5 permits relatively rapid
35 breakdown in the intestine since only a small amount of alkalinity is
required to bring the pH to 5.5. The suspension pH adjustment to 5 ± 0.1 is
not critical. The pH could be adjusted up to 5.4 as may be necessary of a
specific formulation. Although a top spray mode fluid bed apparatus is
preferred, the invention is not limited in this respect, and any suitable spray

coating means, including one with a bottom spray, or a pan type coater, may also be utilized.

Depending on their size, the tablets may be ingested individually or, in another embodiment of the invention, may be filled into dissolvable, hard shell capsules, such as gelatin capsules of varying sizes depending on the dosage of medicament desired. If the tablets are to be encapsulated, a hydrophobic anti-adherent, such as talc, is added (range 0.1 to 4% by weight) to the film coated tablets and blended.

The Examples represent preferred embodiments of the present invention. The following examples further describe the materials and methods used in carrying out the invention and are intended to be for illustrative purposes only, and are not intended to limit the scope or spirit of this invention or the claims in any way. All temperatures are expressed in degrees Centigrade unless otherwise indicated and all mesh sizes are U.S. standard ASTM.

EXAMPLE 1

A ddl formulation for 50mg tablets having the following composition was prepared as described below.

COMPOSITION	WEIGHT % OF COMPONENT	WEIGHT % OF FINAL FORMULATION
<u>A: TABLET CORE</u>		
ddl	71.4	65.763
Microcrystalline Cellulose	24.3	22.359
Na Starch Glycolate	3.0	2.762
Magnesium Stearate	1.3	1.184
<u>B: COATING</u> (Based on 8% coat)		
Eudragit L-30-D 55 (dry basis)	87	6.892
Diethyl Phthalate	13	1.039
(pH adjusted to 5.0 ± 0.1)		

The preparation of ddl tablets was commenced by adding ddl, microcrystalline cellulose, sodium starch glycolate and a first portion of magnesium stearate for compaction, into a tumbling type blender. The ingredients were blended for 10 ± 2 minutes. Prior to blending, any of the initial ingredients which were lumpy, were passed through a #20 mesh screen.

The blended mixture was then passed through a #40 size mesh screen and re-placed into the tumbling type blender and blended again for 10 ± 2 minutes. The resulting blend was then slugged using a 3/4" flat face punches. The slugs were then passed through #10 and #20 size mesh
5 screens for sizing.

A second portion of magnesium stearate needed for tableting was then calculated and placed into the tumbling type blender with the granulation prepared for the sizing of the slugs and blended for 10 ± 2 minutes. The resulting blend was then tableted to obtain the desired tablet
10 weight and hardness.

To prepare sufficient quantities of film coating suspension to coat the tablets, Eudragit L-30-D 55 was filtered through a #60 mesh screen to remove any lumps present therein. The filtered Eudragit was weighed and then added with stirring to a tarred vessel containing one-half the amount of
15 water required. The mixture was continuously stirred for 5 minutes or until a uniform mixture was visually evident. With continuous stirring, diethyl phthalate was added to the vessel and stirring continued for 20 minutes or until a uniform mixture was visually evident. A pH meter was then
20 standardized using pH 4 and pH 7 buffers. With continued stirring, a NaOH solution was added to the vessel until a pH of 5.0 ± 0.1 was obtained. The formula weight of the coating suspension was adjusted using water and stirring was continued for an additional 10 minutes.

The tablet coating procedure utilized a fluid bed apparatus with a top spray mode and appropriate distribution plate to allow fluidization of the
25 product (tablet) in the center.

Before coating, the tablets were pre-warmed in the coating unit to a temperature between 45 -50 °C. An inlet temperature of 50 ± 2 °C was determined to be adequate. The spray rate was adjusted to allow uniform coating and adequate drying of the coat. An $8 \pm 0.5\%$ weight gain due to the
30 film coat was determined to be sufficient. After coating, the tablets were dried for approximately 10 minutes at an inlet temperature of approximately 50 °C.

The so formed enteric coated ddl product was found to give excellent protection against gastric acid (at pH of 3) but had excellent
35 release of ddl at pH's above 5.

EXAMPLE 2

A preferred 50 mg ddl formulation in the form of enteric coated tablets was prepared as described below. ddl (50.00 mg), microcrystalline cellulose (17.00 mg) sodium starch glycolate (2.10 mg) and a first portion of magnesium stearate for compaction (0.60 mg) were placed into a suitable tumbling type blender and blended for 10 ± 2 min. Prior to blending, if any of the ingredients required delumping, they were passed through a #20 mesh screen.

The blended mixture was then passed through a #40 size mesh screen and re-placed into the tumbling type blender and blended again for 10 ± 2 minutes. The resulting blend was then slugged using a 3/4" flat face punches to obtain slugs with a weight and hardness of 1 ± 0.2 g and 15 - 20 SCU, respectively. The slugs were then passed through #10 and #20 size mesh screens.

A second portion of magnesium stearate (0.3 g) for tableting was then placed into the tumbling type blender with the slugs and blended for 10 ± 2 minutes. The resulting blend was then tableted using 3/16" round, plain, deep concave punches to the desired weight at a tablet hardness of 3-6 SCU.

A quantity of film coating suspension in an amount (g) per 100 g for coating the 50 mg ddl tablets was commenced by adding approximately 50 g of water into a suitable container having a stirring mechanism. While the water was being stirred moderately, 33.33 g Eudragit L 30 D-55 was slowly added thereto. Before adding to the water, the Eudragit L 30 D-55 was screened through a #60 size mesh screen.

With continued stirring, 1.50 g diethyl phthalate was added to the water/Eudragit mixture until the diethyl phthalate was in complete solution. While stirring, a sufficient quantity of sodium hydroxide solution (0.1 to 1N) was slowly added in order to adjust the pH of the suspension to 5.0. With continued stirring, water was added to make the formula weight, and the suspension was stirred for an additional 10 minutes.

The tablet coating procedure utilized an Aeromatic Table-top (STREA-1) fluid bed apparatus, with top spray mode and appropriate distribution plate to allow fluidization of the product (tablet) in the center. The coating conditions utilized in the process were:

Charge	250 g
Heat Setting	60 °C
Fan Setting	14

	Inlet Temperature	50 °C
	Pre-heating Time	5 min.
	Spray rate, first 5 min.	4 g/min.
	Spray rate, final	8 g/min.
5	Nozzle opening	1.1 mm
	Air Volume	120
	Outlet Temperature	36 °C
	Weight gain	8%
10	Final drying at fan setting 10	10 min.

Before coating, the tablets were pre-warmed in the coating unit to a temperature between 45 - 50 °C. An inlet temperature of 50 ± 2 °C was determined to be adequate. The spray rate was adjusted to allow uniform coating and adequate drying of the coat. An $8 \pm 0.5\%$ weight gain due to the film coat was determined to be sufficient. After coating, the tablets were dried for approximately 10 minutes at an inlet temperature of approximately 50 °C.

The so formed enteric coated ddl product was found to provide excellent protection against gastric acid (at pH of 3) but had excellent release of ddl at pH's above 5.

Claims

We claim:

- 5
1. An enteric coated pharmaceutical composition comprising a core in the form of a tablet and having an enteric coating surrounding said core, said core comprising an acid labile medicament, a binder or filler, a disintegrant, and a lubricant, said enteric coating comprising a methacrylic
10 acid copolymer, and a plasticizer, and imparting protection to said core so that said core is afforded protection in a low pH environment of 3 or less while capable of releasing medicament at a pH of 4.5 or higher.
 2. The pharmaceutical composition according to claim 1, wherein
15 the pH of said acidic enteric coating is adjusted using alkalizing agents to improve stability between said acid labile medicament in said core and said acidic enteric coating.
 3. The pharmaceutical composition according to claim 2, wherein
20 adjustment of the pH of said acidic enteric coating eliminates incompatibility between said acid labile medicament in said core and said acidic enteric coating.
 4. The pharmaceutical composition according to claim 3,
25 wherein adjustment of said pH of said acidic enteric coating eliminates the need for a protective subcoat between said acid labile medicament in said core and said acidic enteric coating.
 5. The pharmaceutical composition according to claim 4, wherein
30 elimination of said protective subcoat permits quicker release of said acid labile medicament in said core.
 6. The pharmaceutical composition according to claim 1, wherein
35 said medicament is ddl.

7. The pharmaceutical composition according to claim 1, wherein said medicament is pravastatin, erythromycin, digoxin, pancreatin, ddA or ddC.

5 8. The pharmaceutical composition according to claim 1, wherein said binder is microcrystalline cellulose, sodium carboxymethylcellulose, Avicel™ PH101, Avicel™ RC 591, Avicel™ CL-611, Ceolus™, ProSolv™ Methocel™ E-5, Starch 1500, Hydroxypropyl Methylcellulose, Polyvinylpyrrolidone, Potassium Alginate or Sodium Alginate.

10

9. The pharmaceutical composition according to claim 8, wherein said binder is microcrystalline cellulose.

10. The pharmaceutical composition according to claim 1, wherein
15 said lubricant is magnesium stearate.

11. The pharmaceutical composition according to claim 1, wherein said disintegrant is sodium starch glycolate, croscarmellose sodium, corn starch, or cross linked polyvinylpyrrolidone.

20

12. The pharmaceutical composition according to claim 11, wherein said disintegrant is sodium starch glycolate.

13. The pharmaceutical composition according to claim 1, wherein
25 said enteric coating includes methacrylic acid copolymer and a plasticizer.

14. The pharmaceutical composition according to claim 13, wherein said methacrylic acid copolymer is Eudragit L-30-D 55.

30 15. The pharmaceutical composition according to claim 13, wherein said plasticizer is diethyl phthalate, triethyl citrate, triacetin, tributyl sebacate, or polyethylene glycol.

16. The pharmaceutical composition according to claim 15,
35 wherein said plasticizer is diethyl phthalate.

17. The pharmaceutical composition according to claim 1, having the following composition:

5	Material	% (range)
	<u>CORE</u>	
	Drug (didanosine)	1 - 100
	Microcrystalline Cellulose	0 - 40
	Na Starch Glycolate	0 - 6
10	Magnesium Stearate	0 - 3
	<u>COATING</u>	
	Eudragit L-30-D 55	2 - 30
	Diethyl Phthalate	0.5 - 6.0
15		

18. The pharmaceutical composition according to claim 1, wherein said tablet comprises ddl in an amount sufficient to attain a dosage for twice daily administration.

20

19. The pharmaceutical composition according to claim 1, wherein said tablet comprises ddl in an amount sufficient to attain a dosage for once daily administration.

20. An enteric coated ddl comprising a core in the form of a tablet which includes ddl in an amount within the range of from about 1 to about 99.5% by weight of said core, and an enteric coating which includes a methacrylic acid copolymer.

21. ddl as defined to claim 20, in the form of tablets.

22. ddl as defined in claim 21, wherein the enteric coating includes a methacrylic acid copolymer and a plasticizer.

23. ddl as defined in claim 21, wherein said tablet comprises ddl in an amount sufficient to attain a dosage for twice daily administration.

24. The pharmaceutical composition according to claim 21, wherein said tablet comprises ddl in an amount sufficient to attain a dosage for once daily administration.

25. A process for the preparation of an enteric-coated pharmaceutical composition comprising the steps of:

- (a) preparing a blend suitable for tableting;
- 5 (b) tableting said blend into a plurality of individual tablets; and
- (c) coating said tablets.

26. The process according to claim 25, wherein said preparing step (a) comprises the steps of

- 10 (a) blending a mixture comprised of a medicament, a binder, a disintegrant, and a first portion of a lubricant;
- (b) passing said mixture through a #40 size mesh screen;
- (c) reblending said mixture;
- (d) slugging said blend and passing the formed slugs
- 15 successively through #10 and 20 size mesh screens; and
- (e) adding a second portion of lubricant and re-blending.

27. The process according to claim 26, wherein said mixture formed in step (c) is slugged using a 3/4" flat face punch.

20

28. The process according to claim 25, wherein said tableting step is performed using 3/16" round, plain, deep concave punches to obtain tablets of the desired weight.

25 29. The process according to claim 25, wherein said tablets have a hardness of 3-6 SCU.

30. The process according to claim 25, wherein preparation of said coating comprises the steps of:

- 30 (a) mixing a methacrylic acid copolymer with approximately half the amount of water required to attain the desired formulation;
- (b) adding a plasticizer to the mixture of step (a) and stirring until said plasticizer is in complete solution;
- (c) adding sodium hydroxide solution to adjust the pH of the
- 35 mixture of step (b) to 5.0; and
- (d) while stirring the mixture of step (c) adding water in a sufficient amount to bring the mixture to its desired formula weight.

31. The process according to claim 25, wherein said coating step (c) comprises the steps of:

- 5 (a) pre-warming said tablets in a fluid bed spraying apparatus to approximately 45 - 50 °C;
(b) spraying said tablets with said coating; and
(c) drying said tablets.

32. The process according to claim 25, wherein said medicament
10 is an acid labile drug.

33. The process according to claim 32, wherein said medicament is ddl.

15 34. The process according to claim 26, wherein said binder is microcrystalline cellulose.

35. The process according to claim 26, wherein said lubricant is magnesium stearate.
20

36. The process according to claim 26, wherein said disintegrant is sodium starch glycolate.

37. The process according to claim 25, wherein said enteric
25 coating includes methacrylic acid copolymer and diethyl phthalate.

38. The process according to claim 37, wherein said plasticizer is diethyl phthalate.

30 39. The process according to claim 37, wherein said methacrylic acid polymer is Eudragit L-30-D 55.

40. A process for the preparation of an enteric-coated pharmaceutical composition comprising the steps of:
35 (a) preparing a wet mass blend by a wet granulation process for the formation of granules for tableting;
(b) tableting said granules into a plurality of individual tablets;

- (c) coating said tablets with an enteric coating polymer and plasticizer in an aqueous media; and
- (d) blending said coated tablets with an anti-adherent.

5 41. The process according to claim 40, wherein said preparing step (a) comprises the steps of:

- (a) blending a mixture comprised of a medicament, a binder, a disintegrant, and a filler;
- 10 (b) adding a predetermined amount of water or granulation solvent to form a wet blend;
- (c) sizing said blend into granules to aid drying;
- (d) drying said granules to remove excess moisture;
- (e) sizing said dried granules into granules suitable for tableting;
- 15 (f) adding lubricant, disintegrant, one or more fillers, one or more binders, and other excipients for tableting said granules.

 42. The process according to claim 40, wherein preparation of said coating comprises the steps of:

- 20 (a) mixing a methacrylic acid copolymer with approximately 50 g of water;
- (b) adding a plasticizer to the mixture formed in step (a) and stirring until said plasticizer is in complete solution;
- (c) adding sodium hydroxide solution to adjust the pH of the mixture of step (b) to 5.0; and
- 25 (d) while stirring the mixture of step (c) adding water in a sufficient amount to bring the mixture to its desired formula weight .

 43. The process according to claim 41, wherein said coating step comprises the steps of:

- 30 (a) pre-warming said tablets in a fluid bed spraying apparatus to approximately 45 - 50 °C;
- (b) spraying said tablets with said coating; and
- (c) drying said tablets.

35 44. The process according to claim 41, wherein said medicament is an acid labile drug.

45. The process according to claim 44, wherein said medicament is ddl.

46. The process according to claim 41, wherein said binder is microcrystalline cellulose.

47. The process according to claim 41, wherein said lubricant is magnesium stearate.

48. The process according to claim 41, wherein said disintegrant is sodium starch glycolate.

49. The process according to claim 40, wherein said enteric coating includes methacrylic acid copolymer and diethyl phthalate.

50. The process according to claim 49, wherein said plasticizer is diethyl phthalate.

51. The process according to claim 49, wherein said methacrylic acid polymer is Eudragit L-30-D 55.


52. The process according to claim 40, wherein said anti-adherent is talc.

53. The process according to claim 40, further comprising the step of filling said coated tablets prepared in step (d) into a capsule.

30

INTERNATIONAL SEARCH REPORT

 International application No.
 PCT/US98/16323

A. CLASSIFICATION OF SUBJECT MATTER IPC(6) : A61K 9/16, 9/54, 9/62 US CL : 424/451, 457, 458, 459, 461, 490, 489, 494, 497 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) U.S. : 424/451, 457, 458, 459, 461, 490, 489, 494, 497 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5,510,114 A (BORELLA et al.) 23 April 1996, abstract; column 3, lines 1-65 column 4, lines 1-33; columns 5-8.	1-5, 8-20, 25-32, 34-44, 46-53
Y	US 5,422,121 A (LEHMANN et al.) 06 June 1995, abstract; column 7, lines 53-68; column 8, lines 1-12.	31, 43
Y	US 5,556,839 A (GREENE et al.) 17 September 1996, abstract; column 11, lines 37-67; column 12 lines 1-49.	1, 7-17, 25-30, 34-42, 46-52
Y	US 5,326,570 A (RUDNIC et al.) 05 July 1994, abstract; column 2, lines 18-30; column 3, lines 10-68; column 4, lines 1-50; columns 5-7.	1, 7-17, 25-30, 34-42, 46-53
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
A	document defining the general state of the art which is not considered to be of particular relevance	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
E	earlier document published on or after the international filing date	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
L	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
O	document referring to an oral disclosure, use, exhibition or other means	*G* document member of the same patent family
P	document published prior to the international filing date but later than the priority date claimed	
Date of the actual completion of the international search 25 SEPTEMBER 1998		Date of mailing of the international search report 21 OCT 1998
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230		Authorized officer BRIAN K. SEIDLECK  Telephone No. (703) 308-1235